



Reservoir of the European chestnut diversity in Switzerland

S. Pereira-Lorenzo¹ · Y. Bischofberger² · M. Conedera³ · P. Piattini⁴ ·
J. Crovadore⁵ · R. Chablais⁵ · A. Rudow^{6,7} · S. Hatt⁷ · A. M. Ramos-Cabrer¹ ·
T. Barreneche⁸ · F. Lefort⁵

Received: 5 September 2019 / Revised: 16 March 2020 / Accepted: 30 March 2020 /

Published online: 6 April 2020

© Springer Nature B.V. 2020

Abstract

In Switzerland, chestnut forests cover about 27,100 ha, plus some 6800 ha of mixed stands. Due to environmental and historical reasons, most of these still existing forests are located in the Swiss Southern Alps, whereas in the northern parts of the country the chestnut cultivation and the related knowledge strongly regressed since the Little Ice Age period. Nevertheless, Switzerland still hosts valuable genetic resources of the sweet chestnut tree. The present genetic study bases on a nationwide inventory, identification and precise localisation of old and/or grafted chestnut trees for conservation purposes. The main objectives were: (1) to evaluate the genetic diversity and the genetic structure of *Castanea sativa* in Switzerland, and (2) to define a program of conservation including the proposal of a defined core collection. We genetically analysed a pre-selection of 962 accessions (out of 14,165 inventoried trees throughout Switzerland), profiling them with 24 microsatellites. We identified 675 different genotypes out of 962 accessions with a 29.8% of repetitiveness due to clonality. A structural analysis based on a Bayesian method allowed to identify two main clusters, one mostly related to the genetic group from southern Europe (Reconstructed Panmictic Population RPP1) and a second one (RPP2) which revealed to be independent and genetically different from other European groups of chestnut cultivars. The Swiss RPP2 represents a new genetic group, and consequently a complement to genetic resources of chestnut tree in Europe. Genetic analysis allowed defining a core collection of 46 genotypes, which should be used in priority for the Swiss conservation program.

Keywords European chestnut · SSRs · Genetic diversity · Structure · Core collection

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10531-020-01970-2>) contains supplementary material, which is available to authorized users.

Communicated by Daniel Sanchez Mata.

This article belongs to the Topical Collection: Forest and plantation biodiversity.

✉ S. Pereira-Lorenzo
santiago.pereira.lorenzo@usc.es

Extended author information available on the last page of the article

Introduction

Paleobotanical studies on fossil data allowed to identify the main refugia of the Sweet chestnut (*Castanea sativa*) during the Last Glacial Maximum (ca. 20 ka ago) in Transcaucasia and the Italian and Iberian peninsulas, with some minor other spots in the Marmara and Levant regions, in south-eastern Balkans and Central France (Krebs et al. 2019). The post-glacial natural dispersion during the climatic warming in the first half of the Holocene (ca. 11.5–5.7 ka ago) has been totally masked by the subsequent anthropogenic diffusion and cultivation of the species (Conedera et al. 2004). In Western Europe, this happened during the Roman expansion in particular, when the Conqueror diffused the idea of cultivating the chestnut tree outside the Mediterranean regions. As a result, the present genetic structure of the chestnut tree in Europe displays a differentiation between the western Italian and Iberian populations and the eastern Greek and Turkish ones (Mattioni et al. 2013).

Romans privileged the chestnut cultivation as coppice system for timber production, but they already knew the grafting techniques and also reported on existing chestnut varieties (Conedera et al. 2004). The use of chestnut trees for fruit production raised in the post-Roman time and during the Medieval Warm Period in particular (MWP, defined in here as AD 900–1300, Bradley et al. 2003), especially among mountain populations in central and southern-Europe, who cultivated fruit chestnut trees for staple food production as well (Pitte 1986; Squatriti 2013; Buonincontri et al. 2015).

Switzerland followed this main stream. The chestnut tree cultivation was first introduced south of the Alps, mostly as coppice wood, concomitantly with the Roman conquest (Conedera et al. 2004). During the Middle Ages, this cultivation turned to an autarchic *chestnut civilisation* where the chestnut fruit was the main source of food (Krebs et al. 2014). In the northern slope and the inner alpine valleys of the Swiss Alps, the sweet chestnut started to expand in the medieval warm period and was limited to spots in suitable sites in terms of both mild climate and non-calcareous bedrock (Fig. 1; Tanner 1927; Closuit 1958; Furrer 1958, 1972; Heiniger and Conedera 1994).

North of the Alps the chestnut culture was abandoned with the first climatic turbulences in the seventeenth century related to the “Little Ice Age” that may have severely restricted the chances of a normal fruit ripening (Engler 1901). Contrarily, the southern Swiss chestnut civilization survived the climatic reverses and only slowly declined as a consequence of the introduction of alternative staple food (potato and maize), the onset of old and new pathogens (*Phytophthora* spp.; *Cryphonectria parasitica*), and the post-WWII socio-economic changes, which led to an abandonment of traditional chestnut cultivation (Krebs et al. 2014). The arrival of the Asian Gall Wasp (*Dryocosmus kuriphilus*) in 2007 (Forster et al. 2009), repeated summer droughts (Conedera et al. 2010), and the recently emerged fungal pathogen *Gnomoniopsis smithogilvyi* (Pasche et al. 2016) represent additional threats to the Swiss chestnut culture and to restoration efforts.

Nowadays chestnut forests cover about 27,100 ha, to which 6800 ha of mixed stands with at least 50% of chestnut may be added (Brändli 1998). Due to the natural evolution following management abandonment (Conedera et al. 2000), the chestnut area still presenting an orchard structure dropped to 3400 ha. Most of these chestnut orchards are located south of the Alps, where the heritage of the medieval chestnut culture is still present in terms of both living giant trees (Krebs et al. 2012) and knowledge about chestnut tree varieties and related names (Conedera et al. 1994; Gobbin et al. 2007). In the Swiss Northern Alps, on contrary, the cultural heritage including cultivar names almost




Fig. 1 NAP projects to inventory chestnut trees throughout Switzerland

completely extinguished. A few variety names are still in use on the southern bank of Lake Geneva and at least one of them (the variety “Ente”) can be related to accessions located in the Swiss Chablais region. Based on these observations, we can however assume that the whole Switzerland still hosts the historical botanical heritage and the related genetic resources of a medieval sweet chestnut culture.

In order to preserve this heritage, several projects have been carried out in the different cultural and linguistic regions of Switzerland in the context of the National Action Program (NAP) for the conservation and use of plant genetic resources for nutrition and agriculture, established in 1999 and coordinated by the Swiss Federal Office for Agriculture (FOA). NAP represents the Swiss implementation of the Global Plan for the Conservation and Sustainable Utilization of Plant Genetic Resources for Food and Agriculture launched by the Food and Agriculture Organisation (FAO) of the United Nations.

The main objectives of the Swiss chestnut NAP projects were: (1) to localize and inventory the fruit chestnut (*Castanea sativa*) trees in Switzerland; (2) to characterize the identified accessions by a unified methodology; (3) to select the most interesting accessions in terms of phenotypical, morphological and genetic diversity, and (4) to define a core collection to be considered for a conservation program. In this paper, we report on the implementation of the objectives 3 and 4.

The aim of confirming the in situ ethnobotanical and morphological observations with genetic profiling has been implemented from 2010 on, with the additional purpose to avoid redundancies and related costs when setting and maintaining the orchards of the conservation programs.

In this paper, we present the genetic analyses of this nation-wide chestnut accessions inventory campaign that bases on the same set of molecular markers of the EU chestnut database (Pereira-Lorenzo et al. 2017). The obtained results in terms of genetic structure

and diversity are than discussed in the frame of the chestnut cultural heritage at European level.

Material and methods

Project structure, tree identification and localization, and sampling design

Due to the heterogeneity in the history and the related still existing knowledge on the chestnut varieties, the preliminary ethnobotanical, descriptive, and tree inventorial activities were split into different subprojects for each Swiss main cultural area (i.e., the Italian-speaking, southern part, the western French-speaking part and the central-eastern German-speaking part, see Fig. 1 for details).

The NAP projects in southern Switzerland lasted from 1999 to 2018, provided a list of more than one hundred historical, and introduced chestnut variety names (Conedera et al. 1994). For most of the mentioned varieties, we localized and described representative trees in situ (Rudow and Conedera 2001), whereas for a subset of the main varieties, we provided a genotypic analysis and a conservation program in an ex-situ core collection (Gobbin et al. 2007).

Starting in 2011, the development of a comprehensive chestnut traits descriptor's catalogue (Rudow et al. 2012) helped to detect, in the central and eastern parts of northern Switzerland (Fig. 1), an unexpected number and an astonishing diversity of morphologically different tree individuals, which potentially represent ancient varieties of the former local chestnut culture. Finally, we implemented the same approach between 2011 and 2015 also in the western part of northern Switzerland in the frame of a separated initiative.

In 2015, we sampled a subset of 962 accessions (out of the 14,165 inventoried trees throughout Switzerland) for genetic analysis.

Sample selection followed different criteria according to the chestnut areas, for both wild and cultivated trees, and for these latter ones based on the interviews with growers. The presence of grafted trees is confirmed by repetitiveness of the genotypes (Pereira-Lorenzo et al. 2019) and the comparison with the EU database (Pereira-Lorenzo et al. 2017). North to the Alps, emphasis was set on still existing grafted trees, trees based on the visual identification of the grafting marks and/or the information provided by growers, as well as non-grafted old specimens. In southern Switzerland on contrary, many accessions carrying similar names or presenting morphological similitudes were considered for the analysis in order to detect synonyms and homonyms in the inventory. Therefore, several varieties have been considered with redundancies in the southern samples. The final sample consisted of 375 accessions from the western part of northern Switzerland (out of 2255 identified and observed accessions, i.e. about 13%), 385 from the central and eastern part of northern Switzerland (out of 11,390 identified and observed accessions, i.e. about 2.8%) and 202 accessions (out of 520 accessions) from southern Switzerland.

Genetic analysis

DNA extraction

Nucleic acids were extracted from silica gel-dried and apparently healthy portions (50–70 mg) of the leaves sampled in the field using a modified CTAB method adapted from Lefort & Douglas (1999), without any additional treatment of the leaf material. DNA

was resuspended in sterile ultrapure water and quality and concentration were assessed with a Nanodrop ND-1000 spectrophotometer (WITEC AG, Switzerland). All samples were diluted at a final concentration of 50 ng/ μ l.

PCR amplification and SSR analyses

A set of 24 SSRs (Supplementary Table S1) was used for genetic characterization. This set included ten SSR loci provided by Marinoni et al. (2003): *CsCAT1*, *CsCAT2*, *CsCAT3*, *CsCAT6*, *CsCAT8*, *CsCAT14*, *CsCAT15*, *CsCAT16*, *CsCAT17*, and *CsCAT41b*; six other loci described by Buck et al. (2003): *EMCs2*, *EMCs14*, *EMCs15*, *EMCs22*, *EMCs25* and *EMCs38*; four SSR loci by Gobbin et al. (2007): *CIO*, *OAL*, *OCI*, and *RIC*; two SSR loci by Kampfer et al. (1998): *QrZAG96* and *QrZAG4*; and finally two loci described by Steinkellner et al. (1997): *QpZAG36* and *QpZAG110*. The loci *CsCAT16* and *EMCs2* are in the same linkage group, but at a genetic distance over 50 cM, and are therefore considered as independent.

The forward SSR primers were labelled with NED, VIC, 6-FAM or PET fluorochrome, and six different multiplex PCR reactions were designed (Supplementary Table S1).

Data analysis

The presence of null alleles was determined using Micro-Checker ver. 2.2.3 (Van Oosterhout et al. 2004). The observed heterozygosity (H_o) and expected heterozygosity (H_e) (Nei 1978) were estimated using the GenAlEx 6 (Peakall and Smouse 2006). We used the Genodive software package (Meirmans and Van Tienderen 2004) to estimate the inbreeding coefficient (F_{IS}) (Weir and Cockerham 1984).

In order to study population structure and assign individuals to populations based on the SSR genotypes with two alleles, we used a model-based Bayesian procedure, implemented using the STRUCTURE software (Pritchard et al. 2000). Using the admixture model with unlinked loci and correlated allele frequencies, we computed K (unknown) RPPs (reconstructed panmictic populations) of individuals testing $K = 1$ to 15, assuming that the sampled cultivars were from anonymous trees of unknown origin (we used the options $usepopinfo = 0$, $popflag = 0$). This clustering approach assigns individuals probabilistically to reconstructed populations (RPPs) based on genotype. Assignment of a cultivar to a RPP was based on a probability of membership qI of 80%, while a lower probability meant that this accession could have several parental RPPs. Thirty replicate runs per K value were carried out (Porrás-Hurtado et al. 2013), each consisting of a burning period length of 200,000 steps followed by 200,000 MCMC (Monte Carlo Markov Chain) replicates. The STRUCTURE software estimates the most likely number of clusters (K) by calculating the log probability of data for each value of K. According to Evanno et al. (2005), we calculated the second-order change of the likelihood function divided by the standard deviation of the likelihood (ΔK) in order to assess the best K-value supported by the data by using Structure Harvester (Earl and Vonholdt 2012).

PCoA (also known as Classical Multidimensional Scaling) was used with Bayesian-based method (Structure software) to investigate the pattern of population structure. PCoA is a distance-based model which uses jointly a dissimilarity matrix calculated with a simple-matching index, and a factorial analysis. PCoA was performed using the software DARwin 6.0.010 (Dissimilarity Analysis and Representation for Windows). This software

produces graphical representations on Euclidean plans, which preserve at best the distances between units (Perrier et al. 2003; Perrier and Jacquemoud-Collet 2006).

A core collection was defined using PowerCore v. 1.0, which applies the advanced M strategy using heuristic search for establishing core or allele mining sets in large databases, which could represent all alleles (Kim et al. 2007). Three core collections have been calculated, the first one without considering any preselection, the second one considering as pre-selections all those trees retained for conservation, and a third one including the minimum number of trees out of the preselected ones for its maintenance and the necessity to conserve all the allelic diversity.

Results

We identified 675 different genotypes out of 962 accessions evaluated, with 29.8% repetitiveness due to clonality. We found out 54 genotypes repeated two up to 33 times, which totally represented 341 accessions (Supplementary Table S2). All genotypes were checked with the European chestnut database (Pereira-Lorenzo et al. 2017) in the same laboratory and no matches were found.

Marker summary

This study yielded 214 alleles over 24 loci (Table 1) with an average allele number of 8.9 alleles/locus. The most polymorphic loci were *CsCAT3* with 27 alleles, *CsCAT2* with 18 alleles, *EMCs38* with 16 and *CsCAT6* with 14, respectively. The least polymorphic loci were *QrZAG4* and *EMCs25* with 2 alleles and *EMCs2* and *EMCs14* with 3, respectively.

Rare alleles (allele frequency < 0.05) numbered 117 out of 214 total alleles (54.7%, Table 1). No rare alleles were detected in *EMCs25*, *QrZAG4*, and *EMCs2*, and only one was found for *EMCs15*, *RIC* and *CIO*. The maximum number of rare alleles was 22 in *CsCAT3*, followed by the locus *CsCAT2* with 11 rare alleles.

Observed heterozygosity (H_o) varied from 0.024 for *EMCs25* to 0.902 for *CsCAT17* with an average value of 0.578 (Table 1). The lowest expected heterozygosity (H_e) was 0.035 for *EMCs14* and the maximum H_e was 0.887 for *CsCAT2* (Table 1), with an average of 0.648.

The presence of null alleles was detected in the loci *CsCAT14*, *QrZAG4*, *CsCAT2*, *CsCAT41b*, *CIO* and *EMCs25* (Supplementary Table S1).

Genetic and geographic structure

A Bayesian analysis with the STRUCTURE software was conducted using 18 SSRs to determine the genetic structure among 675 unique genotypes. Six loci harbouring null alleles were not included in this analysis. The $\ln[\text{Pr}(X/K)]$ values increased until $K = 2$ (Supplementary Figure S1) estimated by using Structure Harvester (Earl & Vonholdt, 2012) in a group of 400 genotypes out of 675, with a $qI > 80\%$ (59.3% of all genotypes.). This corresponded to a strong differentiation in two main groups of genotypes (RPP, reconstructed panmictic populations), one with 206 genotypes (RPP1), including Swiss cultivars such as ‘Terematt’ (30.5% of the total number of genotypes), and a second one with 194 genotypes (RPP2, 28.7%) including the cultivar ‘Lüina’ (Supplementary Table S2). ‘Terematt’ and ‘Lüina’ have been selected as representative cultivars for RPP1 and RPP2, respectively, because of their broad regional presence in the main Swiss

Table 1 Allelic range (bp), number of alleles per locus (Na), number of rare alleles ($p < 0.05$), percentage of infrequent alleles with respect to the total number of alleles, percentage of infrequent alleles with respect to the total number in each RPP, number of effective alleles (Ne), observed heterozygosity (Ho), expected heterozygosity (He), and alleles identified in the overall set

Locus	bp	Number of alleles(Na)	Rare Alleles ($p < 0.05$)	% of rare alleles with respect to the total number of alleles	Ne	Shannon's Information Index	Ho without hybrids	He without hybrids	He	Fixation Index	Alleles†
<i>CIO</i> [‡]	146–150	5	1	20.0	3.163	1.349	0.418	0.424	0.684	0.388	146, 147, 148, 149, 150
<i>CsCAT1</i>	177–225	11	6	54.5	3.974	1.638	0.757	0.756	0.748	– 0.012	177, 190, 194, 206, 208, 215, 217, 219, 221, 223, 225
<i>CsCAT2</i> [‡]	196–237	18	11	61.1	8.761	2.357	0.509	0.504	0.886	0.425	196, 200 [§] , 205, 209, 211, 213, 215, 217, 219, 221 [§] , 223, 225, 227, 229, 231, 233, 235[§], 237[§]
<i>CsCAT3</i>	190–268	27	22	81.5	7.402	2.455	0.798	0.792	0.865	0.078	190 [§] , 196, 208, 214, 223, 225, 227, 229, 231, 233, 235, 237, 239, 241, 243, 245 [§] , 247, 249, 251, 253, 255, 257, 260, 262, 264, 266[§], 268
<i>CsCAT6</i>	157–196	14	9	64.3	5.536	1.891	0.781	0.782	0.819	0.047	157, 159, 163, 165, 171, 173, 175, 178, 180, 182, 184, 190, 194, 196
<i>CsCAT8</i>	189–212	7	2	28.6	4.309	1.546	0.733	0.732	0.768	0.046	189, 199, 201, 203, 208, 210, 212
<i>CsCAT14</i> [‡]	128–161	6	3	50.0	3.294	1.308	0.688	0.690	0.696	0.012	128[§] , 133, 141, 150, 152, 161
<i>CsCAT15</i>	122–158	9	5	55.6	3.169	1.426	0.696	0.697	0.684	– 0.017	122, 124, 126, 130, 134, 136, 140, 155, 158
<i>CsCAT16</i>	126–156	9	6	66.7	2.777	1.293	0.646	0.641	0.640	– 0.009	126, 130, 132, 136, 141, 143, 145, 148, 156
<i>CsCAT17</i>	129–163	11	4	36.4	6.726	2.070	0.902	0.900	0.851	– 0.059	129, 133, 139, 141, 145, 149, 155, 157, 159[§], 161[§], 163

Table 1 continued

Locus	bp	Number of alleles(Na)	Rare Alleles (p < 0.05)	% of rare alleles with respect to the total number of alleles	Ne	Shannon's Information Index	Ho	Ho without hybrids	He	He without hybrids	Fixation Index	Alleles†
<i>CsCAT41b</i> †	212–235	12	8	66.7	4.515	1.854	0.470	0.469	0.779	0.778	0.396	212, 216, 218, 220, 222, 223, 226, 228, 231, 233, 234, 235
<i>EMCs2</i>	160–166	3	0	0.0	2.465	0.981	0.593	0.594	0.594	0.589	0.003	160, 163, 166
<i>EMCs14</i>	131–144	3	2	66.7	1.036	0.092	0.033	0.028	0.035	0.027	0.067	131, 140, 144
<i>EMCs15</i>	82–93	5	1	20.0	2.782	1.168	0.604	0.608	0.641	0.644	0.057	82, 85, 88, 91, 93
<i>EMCs22</i>	128–155	8	5	62.5	1.613	0.836	0.363	0.372	0.380	0.388	0.046	128, 130, 132, 134, 136, 145, 147, 155§
<i>EMCs25</i> †	158–160	2	0	0.0	1.486	0.509	0.024	0.025	0.327	0.330	0.927	158, 160
<i>EMCs38</i>	232–276	16	9	56.3	7.225	2.227	0.807	0.805	0.862	0.861	0.063	232, 238, 240, 242, 244, 246, 248, 250, 256, 258, 260, 262, 264, 272, 274, 276
<i>OAL</i>	297–332	9	5	55.6	2.639	1.397	0.605	0.602	0.621	0.622	0.026	297, 299§, 301, 303, 305, 309, 322, 330, 332
<i>OCI</i>	146–161	5	2	40.0	2.567	1.104	0.592	0.588	0.610	0.608	0.030	146, 149, 157, 159, 161
<i>QpZAG36</i>	211–225	6	2	33.3	3.649	1.400	0.693	0.694	0.726	0.723	0.045	211, 217, 219, 221, 223, 225
<i>QpZAG110</i>	206–235	13	9	69.2	4.052	1.572	0.662	0.663	0.753	0.752	0.121	206, 210, 213, 215, 219, 220, 222, 223, 225, 227, 229, 233, 235§
<i>QrZAG4</i> †	110–114	2	0	0.0	1.253	0.354	0.195	0.189	0.202	0.198	0.035	110, 114
<i>QrZAG96</i>	137–169	9	4	44.4	3.791	1.535	0.675	0.668	0.736	0.733	0.084	137, 145, 153, 155, 159, 161, 165, 167, 169§
<i>R/C</i>	119–127	4	1	25.0	2.768	1.093	0.631	0.630	0.639	0.638	0.013	119, 121, 123, 127
Total		214	117	54.7								
Mean		8.9			3.790	1.394	0.578	0.577	0.648	0.646	0.117	

Table 1 continued

Locus	bp	Number of Rare Alleles(Na)	Rare Alleles (p < 0.05)	% of rare alleles with respect to the total number of alleles	Ne	Shannon's Information Index	Ho	Ho without hybrids	He	He without hybrids	Fixation Index	Alleles [‡]
SE					0.415	0.121	0.047	0.047	0.044	0.044	0.044	0.044

[†]Locus with null alleles detected with Micro-Checker (van Oosterhout et al. 2004)

[‡]Alleles identified. §, alleles present in Switzerland not detected in previous work in Europe (Pereira-Lorenzo et al. 2010, 2017, 2019); simple underlined, exclusive alleles of the RPP1; double underlined, exclusive alleles of the RPP2; in bold, alleles exclusives of admixed

chestnut area in southern Alps and their charismatic role as a typical early ripening cultivar ('Terematt') and variety to be transformed in dry chestnuts and flower ('Lüina'). This analysis uncovers the first structural level in these data, which corresponds to two main distinguished populations. Geographical distribution of the genotypes for each RPP showed that the two groups were distributed in most of the areas. In many zones of northern Switzerland the two RPPs are both present too with a predominance of RPP2 in the central zone of northern Switzerland, while where RPP1 was predominant in some zones of the western and eastern parts of northern Switzerland Fig. 2). Admixed genotypes were scattered everywhere, even in some valleys with no presence of RPP1 or RPP2.

RPP1 had 14 genotypes repeated two or more times (76 accessions) with a peak of 21 accessions (Fig. 3, Supplementary Table S2); some others as 'Terematt' and 'Fugascera' were found 9 times, 'Marrone dei Pirenei' 6 times, 'Maron Bregaglia', and 'Temporiva' three times each. 'Rosone' was found to be a genetic synonym of 'Belusciora', as 'Roséra' a genetic synonym of 'Temporiva'. RPP2 had 20 repeated genotypes (177 accessions) up to 33 accessions of an unknown variety, 'Magreta Gambarogno' (putative synonym of 'Marrone di Cuneo') was found 3 times, 'Lüina' 29 times, 'Castagno Grande' 21 times, 'Buné Negro' 9 times and 'Berogna' two times. 'Magreta Gambarogno' was revealed to be a genetic synonym of the imported 'Marrone di Cuneo'. We also found repeated genotypes among the admixed genotypes with 'Moréla' repeated 14 times, 'Torcion Negro' and 'Magreta Verzasca' 11 times, 'Tenasca' 5 times and 'Negra' 4 times among others. 'Reveltana' was found to be a genetic synonym of 'Moréla'. In a European context, the Swiss RPP1 genotypes grouped with most of the cultivars compiled in the European database (Pereira-Lorenzo et al. 2017), while RPP2 did not match with any of them (data not shown), making therefore RPP2 a very particular and interesting genotypic group specific to Switzerland, and possibly a remain of the medieval chestnut culture in Switzerland.




Fig. 2 Distribution of 675 Swiss chestnut unique genotypes in reconstructed populations (RPPs) when $K = 2$ based on data of 18 SSR loci




Fig. 3 Location of the main genotypes clonally propagated by RPP when $K = 2$

The geographical distribution of the most repeated genotypes were found mainly in three areas. The most repeated genotype in RPP1 was located only in the South and in the West, meanwhile in northern Alps were only cultivated some genotypes from RPP2 (Fig. 3).

PCoA analysis utilising DARwin 6.0.010 software confirmed the results obtained with STRUCTURE (Fig. 4). Chestnut cultivars grouped in two main clusters differentiated in the first axis, which was consistent with the main RPPs obtained with STRUCTURE with admixed genotypes in between. Additionally, the Neighbor-joining (NJ) tree based on the genetic distance matrix between 673 chestnut genotypes also grouped them in agreement with STRUCTURE for $K = 2$ (Fig. 5) although without any sharp separation.

Pairwise F_{st} value between the two reconstructed main populations RPP1 and RPP2 was 0.096 ($P < 0.001$), while it was and 0.029 ($P < 0.001$) between RPP1 and admixed genotypes and 0.027 ($P < 0.001$) between RPP2 and admixed genotypes, respectively (Supplementary Table S3).

When we compared the genetic diversity for each RPP (Supplementary Table S1), we found 197 alleles in RPP1 and 163 in RPP2. RPP1 displayed 91 rare alleles ($P < 0.05$) while 84 rare alleles were found in RPP2. The total of exclusive alleles, meaning the alleles present in only one RPP, was 25 (11.7%). Most of the exclusive alleles were found in the ‘Terematt’ group (RPP1) with 12 alleles, followed by the admixed group with nine alleles and the ‘Lüina’ group (RPP2) with four alleles.

Among the 24 SSRs analysed in this study, 34 alleles (15.9%) were only found in Switzerland and were not present in the EU database (Pereira-Lorenzo et al. 2017). Moreover, out of these 34 alleles, 12 alleles were exclusive to RPP1, four to RPP2, nine to the admixed genotypes, whereas the remaining nine alleles were found in more than one RPP and/or admixed genotypes. Finally, 36 alleles found in previous studies in southern Europe with the same SSRs did not appear in the Swiss samples, falling to 14 alleles if




Fig. 4 Principal coordinates analysis (PCoA). PCoA using 18 SSRs in the set of 675 Swiss chestnut genotypes showing structure ($K = 2$): in red RPP1 with a $q_i > 80\%$; in blue, RPP2 with a $q_i > 80\%$; in black admixed

hybrids were excluded. We compared those Swiss genotypes with exclusive alleles with previous studies carried out with 10 common SSRs performed in interspecific hybrids (Pereira-Lorenzo et al. 2010, 2019). This showed that 20 Asian alleles were found in 28 Swiss genotypes, but still 14 alleles were only present in 31 Swiss genotypes (Table 1). In order to check if these introgressants could have affected the structure analysis, 28 putative hybrids genotypes were removed: 17 for RPP1, two for RPP2 and nine for the admixed group. The genetic structure without those genotypes harbouring exclusive alleles for Switzerland showed similar results with two RPPs (data not shown).

Core collection as a strategy for conservation

A total of 46 genotypes were selected (Supplementary Table S2) in the first core collection (without considering any preselection) by using heuristic search with PowerCore (v. 1.0) (Kim et al. 2007). This core collection represented 7.1% of the total number of the total 647 genotypes and 4.9% of the total accessions (out of which 28 putative hybrids were excluded). Out of this core collection, only 19 genotypes (11.1%) coincided with trees preselected by the Swiss National Action Plan for conservation (171 genotypes in total and 12 more putative hybrids). When all preselections were considered (171), 188 genotypes

SSR markers and diversity

The number of alleles and average number of alleles per locus was identical to the values referred for the EU chestnut database, even though some differences were found for some loci and the number of genotypes analysed in the present study was much higher. The total expected heterozygosity (H_e) was also similar in both studies, while the total observed heterozygosity (H_o) was found lower in Switzerland, which meant a higher deficit in heterozygosity. A high number of rare alleles (54.7% of all alleles) as well as a high number of exclusive alleles unique to Switzerland (15.9%) give an insight of a particular diversity of chestnut genetic resources in Switzerland.

Genetic and geographical structure and diversification process

The genetic structure revealed by this study was related to two main groups. The first one referred to as ‘Terematt’ (RPP1) belongs to the European group of cultivars listed in the European Database (Pereira-Lorenzo et al. 2017). Surprisingly, this Swiss genetic group did not match with the Italian and French groups as described in the European Database, nor with the genetics of the giant trees from southern Europe (Pereira-Lorenzo et al. 2019). Moreover, the second main cultivar group referred to as ‘Lüina’ (RPP2), did not match with any of the European chestnut cultivars genetic profiles of the EU database, which indicates a new and very different and independent group for southern Europe. These surprising results illustrate that the nowadays remains of the abandoned Swiss chestnut culture might provide a historical window to late medieval chestnut culture in Switzerland, which has been only partly overprinted by more recently varieties, as found in RPP1.

In addition, both Swiss RPPs were related to grafted cultivars and some many other unique putatively wild genotypes in most of the sampled places. As it happened in Spain and Italy, grafted trees in this study clustered with wild trees indicating common ancestors (Pereira-Lorenzo et al. 2019), and, in some cases with genotypes from other areas, such as cv. ‘Marrone di Cuneo’ from Italy, similarly to what happened with the Spanish cultivar ‘Luguesa’ in north-western Spain, which originated from an ancestral population of Italy.

The genetic structure was related to the geographical distribution of the genotypes in Switzerland, even though both genetic groups overlapped in many places as it happened in southern Europe with Italian genotypes, which introgressed in the Iberian Peninsula from at least the seventeenth century.

In chestnut trees, multilocal selection events were identified as the origin of the main cultivars (Pereira-Lorenzo et al. 2010, 2011), likewise to what was observed in olive (Claros et al. 2000; Besnard et al. 2001) where a high diversity of cultivars was found in the same areas of the presence of wild olive trees. As shown by this study, this also seems to be the case in Switzerland, where clonal genotypes were found in both RPPs, one of them genetically different from the European database of cultivars (Pereira-Lorenzo et al. 2017) including the main cv. ‘Lüina’. This cluster also included cv. ‘Magreta Gambarogno’, which was found to be synonym to the putative cv. ‘Marrone di Cuneo’, a northwestern Italian cultivar sampled with this name in Switzerland. The genetic differentiation between Swiss RPPs was found to be even higher, F_{st} 0.096 ($P < 0.001$), than the one between the Iberian and Italian groups, F_{st} 0.068 ($P < 0.001$).

The clonality rate was 29.8%, which is much lower than the clonality rates of 56.5% reported in the EU chestnut database (Pereira Lorenzo et al. 2017) and 51.5% in the study including European giant chestnut trees (Pereira Lorenzo et al. 2019). This low clonality

rate would very likely be related to the sampling approach and in particular the loss of knowledge on certified varieties and related names in the northern part of the country. In addition, the singular genetic group found in Switzerland (RPP2) showed higher clonality (44.7%) than the RPP1 (37.9%) due to clonal accessions in southern Switzerland. This is probably due to better-preserved chestnut culture and related orchards in southern Switzerland. This fact could also be an explanation for the lower allelic diversity found in this singular Swiss RPP2 group displaying 16% less alleles than the RPP1.

The set of markers used in this study has also been very effective in identifying Asian introgressants into local populations, which amounted to 4.1% in this study in comparison to 2.3% of such hybrids in the EU database (Pereira-Lorenzo et al. 2017). Introgressants were present in both RPPs, but more frequent in RPP1. As shown previously without those putative interspecific hybrids, the genetic structure was not affected, which was consistent with previous results in some other areas from Europe and related with the European origin of most of the interspecific hybrids.

Core collection for conservation

In this study, the number of genotypes kept for the core collection without considering any preselection was reduced up to the 92.9% (91.5% if preselections are considered), which was close to the 89% reduction obtained by Kim et al. (2007), and even lower at 95.1% when the total accessions (excluding putative hybrids) were considered. This is in agreement with the values considered by van Hintum et al. (2000), who estimated that a core collection should be no more than 10% of the genotypes, and between 5 and 20% of the accessions of the complete collections. Those obtained values are lower than the ones obtained for the EU database (Pereira-Lorenzo et al. 2017), i.e. 30.0% of the total genotypes (37 cultivars) and 13.9% of the accessions (266 accessions), respectively. These higher values could be partly due to the evaluation of a much wider area at European level.

Conclusions

A new genetic group not reported previously, genetically different from the main European group of cultivars was found in Switzerland and might represent a remain of the abandoned chestnut culture of the Medieval times.

The genetic analysis carried in this study allowed defining a priority conservation program for the Swiss European chestnut trees based on a core collection of 53 genotypes. Without regard to typical regional varieties, the minimal core collection could even be reduced to 46 genotypes, corresponding to 7% of the total genotypes evaluated in this study. This core collection based on allelic diversity could be completed with some other genotypes with phenotypic traits of interest, to minimize the risk of losing relevant genetic diversity.

Acknowledgements This study was funded by the Swiss Federal Office for Agriculture FOAG in the framework of the Swiss National Plan of Action for plant genetic resources for food and agriculture. We thank to the Spanish government for the funds under the contract AGL2013-48017-C2-2-R to compare allelic ranges with Spanish samples.

Author Contributions Designed research: SP-L, YB, TB, FL. Performed research: SP-L, YB, MC, PP, JC, RC, AR, SH, AMR-C, TB, FL. Contributed new reagents or analytical tools: SP-L, MC, AMR-C, HC, TB, FL. Analyzed data: SP-L, MC, RC, AMR-C, TB, FL. Wrote the paper: SP-L, YB, MC, PP, AR, SH, AMR-C, TB, FL.

Data Accessibility Statement A comprehensive DNA samples database is stored at the Institute Land Nature Environment (HES-SO University of applied sciences and arts western Switzerland).

References

- Besnard G, Baradat P, Berville A (2001) Genetic relationships in the olive (*Olea europaea* L.) reflect multilocal selection of cultivars. *Theor Appl Genet* 102:251–258. <https://doi.org/10.1007/s001220051642>
- Bradley RS, Hughes MK, Diaz HF (2003) Climate in medieval time. *Science* 302:404–405. <https://doi.org/10.1126/science.1090372>
- Brändli UB (1998) Die häufigsten Waldbäume der Schweiz. Ergebnisse aus dem Landesforstinventar 1983–85: Verbreitung, Standort und Häufigkeit von 30 Baumarten, 2nd edn. Eidgenössische Forschungsanstalt für Wald, Schnee und Landschaft, Birmensdorf
- Buck EJ, Hadonou M, James CJ, Blakesley D, Russell K (2003) Isolation and characterization of polymorphic microsatellites in European chestnut (*Castanea sativa* Mill.). *Mol Ecol Notes* 3:239–241. <https://doi.org/10.1046/j.1471-8286.2003.00410.x>
- Buonincontri MP, Saracino A, Di Pasquale G (2015) The transition of chestnut (*Castanea sativa* Miller) from timber to fruit tree: Cultural and economic inferences in the Italian peninsula. *Holocene* 25(7):1111–1123. <https://doi.org/10.1177/0959683615580198>
- Claros MG, Crespillo R, Aguilar ML, Canovas FM (2000) DNA fingerprinting and classification of geographically related genotypes of olive-tree (*Olea europaea* L.). *Euphytica* 116:131–142. <https://doi.org/10.1023/A:1004011829274>
- Closuit R (1958) Le châtaignier dans la vallée suisse du Rhône. *Mitt Eidgenöss Anst Forstliche Vers* 34(3):183–220
- Conedera M, Barthold F, Torriani D, Pezzatti GB (2010) Drought sensitivity of *Castanea sativa*: case study of summer 2003 in the southern Alps. *Acta Hortic* 866:297–302
- Conedera M, Krebs P, Tinner W, Pradella M, Torriani D (2004) The cultivation of *Castanea sativa* (Mill.) in Europe from its origin to its diffusion on a continental scale. *Veg Hist Archaeobot* 13:161–179. <https://doi.org/10.1007/s00334-004-0038-7>
- Conedera M, Müller-Starck G, Fineschi S (1994) Genetic characterization of cultivated varieties of European Chestnut (*Castanea sativa* Mill.) in Southern Switzerland I Inventory of chestnut varieties: history and perspectives. In: Proceedings of the International Congress on Chestnut Spoleto Italy 20–23 October 1993, pp 299–302.
- Conedera M, Stanga P, Lischer C, Stöckli V (2000) Competition and dynamics in abandoned chestnut orchards in southern Switzerland. *Ecol Mediterr* 26(1/2):101–112
- Earl DA, Vonholdt BM (2012) STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conserv Genet Resour* 4:359–361. <https://doi.org/10.1007/s12686-011-9548-7>
- Engler A (1901) Über Verbreitung Standortansprüche und Geschichte der *Castanea vesca* Gärtner mit besonderer Berücksichtigung der Schweiz. *Ber Schweiz Bot Ges* 9:23–62
- Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol* 14:2611–2620. <https://doi.org/10.1111/j.1365-294X.2005.02553.x>
- Forster B, Castellazzi T, Colombi L, Fürst E, Marazzi C, Meier F et al (2009) Die Edelkastaniengallwespe *Dryocosmus kuriphilus* (Yasumatsu) (Hymenoptera Cynipidae) tritt erstmals in der Südschweiz auf. *Mitt Schweiz Entomol Ges* 82:271–279. <https://doi.org/10.5169/seals-402995>
- Furrer E (1958) Die Edelkastanie in der Innerschweiz. *Mitt Eidgenöss Anstalt Forstl Vers* 34(3):90–182
- Furrer E (1972) Die früheste Kastanienkultur in der Innerschweiz und die heutigen Reste. *Ber Schweiz Bot Ges* 82(1):5–13
- Gobbin D, Hohl L, Conza L, Jermini M, Gessler C, Conedera M (2007) Microsatellite-based characterization of the *Castanea sativa* cultivar heritage of southern Switzerland. *Genome* 50(12):1089–1103. <https://doi.org/10.1139/G07-086>
- Heiniger U, Conedera M (1994) Chestnut forests and chestnut cultivation in Switzerland. In: Proceedings of the international chestnut conference West Virginia University College of Agriculture and Forestry Morgantown West Virginia July 10–14 1992, pp 175–178
- Kamper S, Lexer C, Glossl J, Steinkellner H (1998) Characterization of (GA)_n microsatellite loci from *Quercus robur*. *Hereditas* 129:183–186. <https://doi.org/10.1111/j.1601-5223.1998.00183.x>

- Krebs P, Koutsias N, Conedera M (2012) Modelling the eco-cultural niche of giant chestnut trees: new insights into land use history in southern Switzerland through distribution analysis of a living heritage. *J Hist Geogr* 38(4):372–386. <https://doi.org/10.1016/j.jhg.2012.01.018>
- Krebs P, Pezzatti GB, Beffa G, Tinner W, Conedera M (2019) Revising the sweet chestnut (*Castanea sativa* Mill.) refugia history of the last glacial period with extended pollen and macrofossil evidence. *Quat Sci Rev* 206:111–128. <https://doi.org/10.1016/j.quascirev.2019.01.002>
- Krebs P, Tinner W, Conedera M (2014) Del castagno e della castanicoltura nelle contrade insubriche: tentativo di una sintesi eco-storica. *Arch Stor Ticin* 155:5–37
- Kim KW, Chung HK, Cho GT, Ma KH, Chandrabalan D, Gwag JG et al (2007) PowerCore: a program applying the advanced M strategy with a heuristic search for establishing core sets. *Bioinformatics* 23(16):2155–2162. <https://doi.org/10.1093/bioinformatics/btm313>
- Lefort F, Douglas GC (1999) An efficient micro-method of DNA isolation from mature leaves of four hardwood tree species *Acer Fraxinus Prunus* and *Quercus*. *Ann For Sci* 56(3):259–263. <https://doi.org/10.1051/forest:19990308>
- Marinoni D, Akkak A, Bounou G, Edwards KJ, Botta R (2003) Development and characterization of microsatellite markers in *Castanea sativa* (Mill). *Mol Breed* 11:127–136. <https://doi.org/10.1023/A:1022456013692>
- Mattioni C, Martín MA, Pollegioni P, Cherubini M, Villani F (2013) Microsatellite markers reveal a strong geographical structure in European populations of *Castanea sativa* (Fagaceae): evidence for multiple glacial refugia. *Am J Bot* 100:951–961. <https://doi.org/10.3732/ajb.1200194>
- Meirmans PG, Van Tienderen PH (2004) Genotype and genodive: two programs for the analysis of genetic diversity of asexual organisms. *Mol Ecol Notes* 4:792–794. <https://doi.org/10.1111/j.1471-8286.2004.00770.x>
- Nei M (1978) Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89:583–590
- Pasche S, Calmin G, Auderset G, Crovadore J, Pelletteret P, Mauch-Mani B et al (2016) *Gnomoniopsis smithogilyvi* prevalence in *Castanea sativa* shoots in Switzerland. *Fungal Genet Biol* 87:9–21. <https://doi.org/10.1016/j.fgb.2016.01.002>
- Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel Population genetic software for teaching and research. *Mol Ecol Notes* 6:288–295. <https://doi.org/10.1111/j.1471-8286.2005.01155.x>
- Pereira-Lorenzo S, Costa R, Ramos-Cabrer AM, Ribeiro C, Serra da Silva C, Manzano G, Barreneche T (2010) Variation in grafted European chestnut and hybrids by microsatellites reveals two main origins in the Iberian Peninsula. *Tree Genet Genomes* 6:701–715. <https://doi.org/10.1007/s11295-010-0285-y>
- Pereira-Lorenzo S, Lourenço Costa RM, Ramos-Cabrer AM, Ciordia-Ara M, Marques Ribeiro CA, Borges O, Barreneche T (2011) Chestnut cultivar diversification process in the Iberian Peninsula Canary Islands and Azores. *Genome* 54:301–315. <https://doi.org/10.1139/g10-122>
- Pereira-Lorenzo S, Ramos Cabrer AM, Barreneche T, Mattioni C, Villani F, Díaz-Hernández MB et al (2017) Database of European chestnut cultivars and definition of a core collection using simple sequence repeats. *Tree Genet Genomes* 13:114. <https://doi.org/10.1007/s11295-017-1197-x>
- Pereira-Lorenzo S, Ramos-Cabrer AM, Barreneche T, Mattioni C, Villani F, Díaz-Hernández MB et al (2019) Instant domestication process of European chestnut cultivars. *Ann Appl Biol* 174(1):74–85. <https://doi.org/10.1111/aab.12474>
- Perrier X, Flori A, Bonnot F (2003) Data analysis methods. In: Hamon P, Seguin M, Perrier X, Glaszmann JC (eds) Genetic diversity of cultivated tropical plants science. Enfield, Science Publishers, Montpellier, pp 43–76
- Perrier X, Jacquemoud-Collet JP (2006) DARwin software. Retrieved from <https://darwin.cirad.fr/>
- Pitte JR (1986) Terre de castanides Hommes et paysages du châtaigniers de l'antiquité à nos jours. Fayard, Paris
- Porrás-Hurtado L, Ruiz Y, Santos C, Phillips C, Carracedo Á, Lareu MV (2013) An overview of STRUCTURE: applications, parameter settings, and supporting software. *Front Genet* 4:98. <https://doi.org/10.3389/fgene.2013.00098>
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* 155(2):945–959
- Rudow A, Bischofberger Y, Piattini P, Hatt S (2012) Handbook of sweet chestnut descriptors (*Castanea sativa* Mill.). Federal Office for Agriculture, Bern
- Rudow A, Conedera M (2001) Blüte und Sortenerkennung bei der Edelkastanie (*Castanea sativa* Mill.) auf der Alpensüdseite der Schweiz. *Bot Helv* 111:1–23
- Squatriti P (2013) Landscape and change in early medieval Italy: chestnuts economy and culture. University Press Cambridge, Cambridge

- Steinkellner H, Fluch S, Turetschek E, Lexer C, Streiff R, Kremer A et al (1997) Identification and characterization of (GA/CT)_n-microsatellite loci from *Quercus petraea*. *Plant Mol Biol* 33:1093–1096. <https://doi.org/10.1023/A:1005736722794>
- Tanner H (1927) Die Verbreitung und wirtschaftliche Bedeutung der zahmen Kastanie im Kanton St Gallen. *Jahrbuch St Gallischen Naturwissenschaftlichen Gesellschaft* 63:27–48
- van Hintum TJJ, Brown AHD, Spillane C, Hodgkin T (2000) Core collections of plant genetic resources IPGRI. (Technical Bulletin No 3). International Plant Genetic Resources Institute, Rome, Italy
- van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Mol Ecol Resour* 4:535–538. <https://doi.org/10.1111/j.1471-8286.2004.00684.x>
- Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. *Evolution* 38:1358–1370. <https://doi.org/10.2307/2408641>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Affiliations

S. Pereira-Lorenzo¹ · Y. Bischofberger² · M. Conedera³ · P. Piattini⁴ · J. Crovadore⁵ · R. Chablais⁵ · A. Rudow^{6,7} · S. Hatt⁷ · A. M. Ramos-Cabrer¹ · T. Barreneche⁸ · F. Lefort⁵

¹ Departamento de Producción Vegetal y Proyectos de Ingeniería, Escola Politécnica Superior, Universidade de Santiago de Compostela, Campus de Lugo, 27002 Lugo, Spain

² In Situ Vivo Sàrl, Chemin de Grésy 6, 1241 Puplinge, Switzerland

³ Swiss Federal Institute for Forest, Snow and Landscape Research WSL, a Ramèl 18, 6593 Cadenazzo, Switzerland

⁴ EcoControl SA, Via Rovedo 16, 6600 Locarno, Switzerland

⁵ Plants and Pathogens Group, Research Institute Land Nature and Environment (inTNE), Hepia, HES-SO University of Applied Sciences and Arts Western Switzerland, Route de Presinge 150, Jussy, 1254 Geneva, Switzerland

⁶ Institute for Terrestrial Ecosystems, Department of Environmental Systems Science, ETH Zurich, 8092 Zürich, Switzerland

⁷ MOGLI Solutions, 5400 Baden, Switzerland

⁸ UMR BFP Biologie du Fruit et Pathologie, INRAE-Université de Bordeaux, F-33882 Villenave d'Ornon, France